

Biophysical and biological
characterization of Huntingtin
interacting protein HYPK

Thesis Submitted for the Degree of Doctor of
Philosophy (Science) of University of Calcutta, India
(2008)

By

Swasti Raychaudhuri

Contents

| | |
|---|-----------|
| Abbreviation | 1 |
| Synopsis | 2 |
| Chapter 1: General Introduction | |
| 1.01 Huntington's disease | 12 |
| 1.02 Neuropathology in HD | 12 |
| 1.03 The Huntingtin (Htt) protein | 13 |
| 1.04 Nuclear Aggregates of Htt: Hallmark of HD | 15 |
| 1.05 Cell death in HD | 17 |
| 1.05.01 Apoptosis in HD | 17 |
| 1.05.02 Htt aggregates and apoptosis | 18 |
| 1.05.03 Other processes impaired in HD | 20 |
| 1.06 Interactors of huntingtin and implication in pathogenesis of HD | 22 |
| 1.07 Transcription deregulation in HD | 25 |
| 1.08 Chaperones: Protein misfolding/refolding, proteosomal degradation and autophagy in HD | 27 |
| 1.09 Intrinsically unstructured proteins in neurodegenerative Diseases | 33 |
| 1.10 Objective of the thesis: Characterization of HYPK, a novel huntingtin interactor | 36 |
| References | 37 |
| Chapter 2: General Materials and Methods | |
| 2.01 Cloning, expression and Purification of HYPK | 52 |
| 2.02 Cloning of N-terminal and C-terminal region of HYPK | 53 |
| 2.03 Cloning of normal and expanded huntingtin exon1 in DsRedC1 vector | 53 |
| 2.04 Cloning of Eukaryotic Translation Elongation Factor 1 α | 53 |
| 2.05 Cloning of HSP70 | 54 |
| 2.06 Cell culture and transfection | 54 |
| 2.07 Mass Spectrometry | 54 |

Chapter 3: Biophysical Characterization of HYPK

| | | |
|----------|--|----|
| 3.01. | Prologue | 56 |
| 3.02 | Protocols for Biophysical Characterization of HYPK | 56 |
| 3.02.1 | Size-exclusion Chromatography | 56 |
| 3.02.2 | Dynamic Light Scattering (DLS) | 56 |
| 3.02.3 | CD Spectroscopy | 57 |
| 3.02.4 | Fluorescence Spectroscopy | 57 |
| 3.02.5 | Fluorescence Spectroscopy of HYPK at Different pH | 57 |
| 3.02.6 | Fluorescence Spectroscopy of HYPK in Presence of Guanidine Hydrochloride and Sodium Perchlorate | 58 |
| 3.02.7 | Acrylamide Quenching Study of HYPK Fluorescence | 58 |
| 3.02.8 | Limited Proteolysis | 58 |
| 3.02.9 | <i>in vitro</i> Cross-linking of HYPK | 59 |
| 3.02.10 | Native Gel Electrophoresis | 59 |
| 3.02.11 | Estimation of Mean Net Charge (R) and Mean Hydrophobicity (H) | 59 |
| 3.02.12 | Analysis of Amino Acid Sequence of HYPK | 60 |
| 3.02.13 | Effect of Ca ⁺⁺ Concentration on HYPK Conformation | 60 |
| 3.03 | Results | 61 |
| 3.03.1 | Expression and purification of recombinant HYPK | 61 |
| 3.03.2 | Unusual Movement of HYPK on SDS Polyacrylamide Gel | 61 |
| 3.03.3 | Size Exclusion Chromatography | 62 |
| 3.03.4 | Conformational Studies of HYPK with Various Spectroscopic Techniques | 65 |
| 3.03.4.1 | Dynamic Light Scattering (DLS) | 65 |
| 3.03.4.2 | Circular Dichroism (CD) | 66 |
| 3.03.4.3 | Fluorescence Spectroscopy | 67 |
| 3.03.5 | Limited Proteolysis | 71 |
| 3.03.6 | Theoretical Analysis of HYPK Sequence | 73 |

| | | |
|---|---|-----|
| 3.03.7 | Effect of Ca ⁺⁺ on HYPK Conformation | 74 |
| 3.04 | Discussions | 79 |
| 3.05 | Conclusion | 87 |
| | References | 88 |
| Chapter 4: Biological Characterization of HYPK | | |
| 4.01 | Prologue | 92 |
| 4.02 | Protocols for Biological Characterization of HYPK | 92 |
| 4.02.1 | Microscopy, Live cell microscopy and Photo-bleaching | 92 |
| 4.02.2 | Co-immunoprecipitation | 94 |
| 4.02.3 | Filter Retardation Assay | 95 |
| 4.02.4 | Detection of Nuclear Fragmentation | 95 |
| 4.02.5 | MTT Assay | 96 |
| 4.02.6 | Fluorometric Determination of Caspase-2, Caspase-3, and Caspase-8 Activity | 96 |
| 4.02.7 | <i>in vitro</i> Chaperone Activity Assays | 97 |
| 4.02.8 | <i>in vivo</i> Chaperone Activity Assay | 97 |
| 4.02.9 | Nascent Peptide Binding Assay | 98 |
| 4.02.10 | Interactors of HYPK other than Htt: Pull Down Assay and Mass Spectrometry | 99 |
| 4.03 | Results | 99 |
| 4.03.1 | Interactions of HYPK with N-terminal Htt coded by exon1 of <i>htt</i> gene | 99 |
| 4.03.2 | Suppression of aggregates formed by N-terminal Htt with 40Q by co-expression of GFP-HYPK | 101 |
| 4.03.3 | Live cell microscopy reveals that co-expression of GFP-HYPK reduces the kinetics of mutated N-terminal Htt mediated aggregate formation | 106 |
| 4.03.4 | Fluorescence recovery after photobleaching (FRAP) and fluorescence loss in photobleaching (FLIP) of Htt aggregates in the presence and | |

| | |
|---|-----|
| absence of GFP-HYPK | 109 |
| 4.03.5 Reduction in SDS Insoluble Aggregates in presence of HYPK | 113 |
| 4.03.6 Reduction of DsRed-H40 induced apoptosis by co-expression of GFP-HYPK | 114 |
| 4.03.7 HYPK has chaperone like activity | 117 |
| 4.03.8 HYPK contains a putative nascent peptide binding motif | 120 |
| 4.03.9 N-terminus and C-terminus of HYPK co-localizes with Htt aggregates when over-expressed | 121 |
| 4.03.10 Experimental validation of interaction of HYPK and nascent peptides | 123 |
| 4.03.11 <i>in vivo</i> chaperone-like activity of C-terminus of HYPK | 125 |
| 4.03.12 HYPK interacts with Eukaryotic Translation Elongation Factor 1 α | 127 |
| 4.04 Discussions | 127 |
| 4.05 Conclusion | 133 |
| References | 134 |
| Chapter 5: Intrinsically Unstructured Proteins in Neurodegenerative diseases | |
| 5.01 Prologue | 137 |
| 5.02 Protocols | 137 |
| 5.02.1 Construction of datasets | 137 |
| 5.02.2 Disorder Prediction | 138 |
| 5.02.3 Unstructuredness and protein interaction | 140 |
| 5.02.4 Functional annotation | 141 |
| 5.02.5 Statistical analysis | 141 |
| 5.03 Results | 142 |
| 5.03.1 Unstructured Proteins are Prevalent in Neurodegenerative Diseases | 142 |
| 5.03.2 Unstructuredness is Prevalent in Hub Proteins in | |

| | |
|--|-------|
| the HD, PD and AD datasets | 144 |
| 5.03.3 Functions of the Unstructured Proteins in Neurodegenerative Diseases (HD, PD and AD) | 146 |
| 5.04 Discussions | 148 |
| References | 150 |
| Chapter 6: Summary and Conclusions | 153 |
| Appendix | i-1ix |