

# **EXPERIMENTAL STUDY OF THE MAGNETIC PROPERTIES OF QUASI-ONE DIMENSIONAL ANTIFERROMAGNETS**

A Thesis Submitted

by

**BHOLANATH PAHARI**

Experimental Condensed Matter Physics Division  
Saha Institute of Nuclear Physics  
1/AF, Bidhannagar, Kolkata-700064, India.

to

**JADAVPUR UNIVERSITY**

Kolkata-700032, India.

For the Degree of  
**Doctor of Philosophy in Science**

**2008**

# Contents

Title Page . . . . .	i
Preface . . . . .	iii
Table of Contents . . . . .	vi
Publications . . . . .	ix
Acknowledgments . . . . .	xi
Dedication . . . . .	xiii
<b>1 Introduction to Quasi-1D Quantum Spin System</b>	<b>1</b>
1.1 Classical Treatments . . . . .	3
1.2 Quantum Mechanical Treatment . . . . .	4
1.2.1 Uniform Exchange $S = 1/2$ HAF Chains . . . . .	5
1.2.2 Alternating Exchange $S = 1/2$ HAF Chain . . . . .	8
1.2.3 Uniform Exchange $S = 1$ HAF chain . . . . .	14
1.3 The Lieb-Shultz-Mattis theorem . . . . .	18
1.4 Effect of Impurity Doping in the Integer-spin chain system . . . . .	21
1.5 Study of quasi-1D HAF system by NMR . . . . .	22
<b>Bibliography</b>	<b>24</b>
<b>2 Basic Theory of Nuclear Magnetic Resonance (NMR)</b>	<b>27</b>
2.1 Basic Resonance Theory . . . . .	27
2.2 Classical Treatment of NMR (Bloch Equations) . . . . .	29
2.3 Broadening of NMR Line . . . . .	32
2.3.1 Natural width due to spontaneous emission . . . . .	32
2.3.2 Width due to Spin-lattice Relaxation . . . . .	32
2.3.3 Broadening due to Field Inhomogeneities . . . . .	32
2.3.4 Other effects . . . . .	33
2.4 Hyperfine Interactions . . . . .	33
2.4.1 Magnetic Interaction . . . . .	34
2.4.2 The Effect of Magnetic Interaction to Resonance Frequency Shift	35
2.4.3 Effect of Magnetic Interaction to Spin-lattice Relaxation . . .	40
2.4.4 Effect of Magnetic Interaction to NMR Line Width . . . . .	40
2.4.5 Electrostatic Interaction . . . . .	42
2.4.6 Effect of Nuclear Quadrupolar Interaction Over Zeeman Interaction	45
2.5 Combined Magnetic and Nuclear Quadrupolar interaction Effect . . .	46

2.6	Powder Pattern of a NMR Line Shape . . . . .	47
2.6.1	Probability distribution function . . . . .	48
2.6.2	NMR line shape . . . . .	49
2.7	Moments of a NMR spectrum . . . . .	49
2.8	Relaxation Processes . . . . .	50
2.8.1	Nuclear Spin-Lattice Relaxation . . . . .	51
2.8.2	Spin-spin relaxation . . . . .	53
2.8.3	Correlation and Nuclear Relaxation Rates . . . . .	54
	<b>Bibliography</b>	<b>56</b>
<b>3</b>	<b>Experimental Methods</b>	<b>59</b>
3.1	Sample Preparation . . . . .	59
3.2	Characterization of Samples by XRD . . . . .	60
3.3	Measurement of Bulk magnetic properties . . . . .	61
3.3.1	SQUID magnetometer: magnetization measurement . . . . .	61
3.3.2	Set-up for heat capacity measurement . . . . .	62
3.4	NMR Measurements . . . . .	65
3.4.1	Pulsed NMR . . . . .	66
3.4.2	Spin-Echo Method . . . . .	67
3.4.3	Dead Time . . . . .	69
3.4.4	Recording of NMR spectra . . . . .	69
3.4.5	Measurement of NMR shift . . . . .	70
3.4.6	Measurement of Spin-lattice Relaxation . . . . .	70
3.4.7	Measurement of Spin-spin Relaxation . . . . .	72
3.4.8	Description of NMR Spectrometer . . . . .	72
	<b>Bibliography</b>	<b>75</b>
<b>4</b>	<b>Experimental Investigations on the Compound BaCu<sub>2</sub>V<sub>2</sub>O<sub>8</sub></b>	<b>76</b>
4.1	Introduction . . . . .	76
4.2	Crystal Structure of BaCu <sub>2</sub> V <sub>2</sub> O <sub>8</sub> . . . . .	77
4.3	Motivations . . . . .	79
4.4	Preparation of the Sample . . . . .	79
4.5	Characterization of the Sample . . . . .	80
4.6	Experimental . . . . .	80
4.7	Results and Discussion . . . . .	81
4.7.1	Magnetic Susceptibility . . . . .	81
4.7.2	NMR Spectra . . . . .	82
4.7.3	NMR Shift . . . . .	85
4.7.4	Nuclear spin-lattice relaxation rates . . . . .	89
4.8	Summary . . . . .	95
	<b>Bibliography</b>	<b>97</b>

<b>5 Experimental Investigations on the Compound SrNi<sub>2</sub>V<sub>2</sub>O<sub>8</sub></b>	<b>100</b>
5.1 Introduction . . . . .	100
5.2 Crystal Structure . . . . .	102
5.3 Outlines of present work . . . . .	102
5.4 Sample Preparation . . . . .	103
5.5 Sample Characterization . . . . .	104
5.6 Experimental . . . . .	105
5.7 Results and Discussions . . . . .	105
5.7.1 Magnetic Susceptibility . . . . .	105
5.7.2 <sup>51</sup> V NMR Spectra . . . . .	107
5.7.3 NMR Shift . . . . .	108
5.7.4 Spin-Lattice Relaxation Rate . . . . .	111
5.7.5 Spin-Spin Relaxation Time . . . . .	112
5.8 Summary . . . . .	114
<b>Bibliography</b>	<b>115</b>
<b>6 Doping Effects in S = 1 Spin Chain Compound SrNi<sub>2</sub>V<sub>2</sub>O<sub>8</sub></b>	<b>117</b>
6.1 Introduction . . . . .	117
6.2 Sample Preparation and Characterization . . . . .	119
6.3 Experimental . . . . .	123
6.4 Experimental Results of Bulk Magnetic Properties Measurements . . . . .	123
6.4.1 Magnetic Susceptibility . . . . .	123
6.4.2 Magnetization . . . . .	129
6.4.3 Heat Capacity and Entropy . . . . .	131
6.4.4 Conclusions of the Results Obtained from Bulk Properties Measurements . . . . .	133
6.5 Results and Discussions of the NMR Measurements . . . . .	134
6.5.1 <sup>51</sup> V NMR SPECTRA . . . . .	134
6.5.2 NMR Shift . . . . .	138
6.5.3 Spin-lattice Relaxation Rate . . . . .	139
6.5.4 Conclusions from NMR Results . . . . .	144
6.6 Summary . . . . .	144
<b>Bibliography</b>	<b>146</b>
<b>7 Conclusions</b>	<b>149</b>
<b>A Application of Dimer Model in BaCu<sub>2</sub>V<sub>2</sub>O<sub>8</sub></b>	<b>152</b>
<b>B Calculation of Magnetic relaxation for I=7/2</b>	<b>155</b>