

**Transcription regulation by Huntingtin
Interacting Protein 1 and its molecular
partner HIPPI: possible role in
Huntington's disease**

**Thesis Submitted for the Degree of Doctor of
Philosophy (Biochemistry) of University of Calcutta,
India
(2011)**

**By
Moumita Datta**

Content

Abbreviations	1
Synopsis	3
Chapter 1: General Introduction	
1.1 Huntington's disease	10
1.2 Genetic and molecular basis of HD	11
1.2.1 Huntingtin mutation	11
1.2.2 The protein- Huntingtin	12
1.2.3 Huntingtin interacting proteins	14
1.2.4 Normal function of Huntingtin	15
1.3 Models of Huntington's disease	18
1.4 Mechanism of neurodegeneration in HD	20
1.4.1 Protein aggregation, dysfunction of ubiquitin proteasome system and chaperone activity	21
1.4.2 Proteolysis of mutant HTT	23
1.4.3 Transcriptional deregulation	23
1.4.4 Mitochondrial dysfunction and oxidative stress	24
1.4.5 Excitotoxicity and striatal dysfunction	25
1.4.6 Apoptosis and autophagy	26
1.4.7 Loss of wild type HTT function	28
1.4.8 Non-cell-autonomous processes in HD	28
1.5 Transcriptional deregulation in HD	30
1.5.1 Early studies on single gene expression	31
1.5.2 Microarray studies	31
1.5.3 Pathways altered in HD: implications from microarray data	35
1.5.4 Mechanism of transcription deregulation in HD	37
1.5.4.1 Aberrant interaction of mutant HTT with transcription factors	37
1.5.4.2 Aberrant interaction of mutant HTT with chromatin modification machinery	39
1.5.4.3 Loss of wild type HTT mediated sequestration of transcription repressor	40
1.5.4.4 Aberrant interaction of mutant HTT with DNA	41
1.5.4.5 Post-transcriptional deregulation by microRNAs	43
1.6 Huntingtin Interacting Protein 1 (HIP1)	44

1.7	HIP1 Protein Interactor (HIPPI)	46
1.7.1	Pro-apoptotic functions of HIPPI: implications from its interacting partners	47
1.7.2	Transcription regulation by HIPPI	47
Chapter 2: Materials and Methods		
2.1	Materials	53
2.1.1	Reagents	53
2.1.2	Mammalian cells	53
2.1.3	Plasmids	53
2.2	Methods	
2.2.1	Gene cloning	55
2.2.2	Mammalian cell culture and transfection	55
2.2.3	Knock down of <i>HIP1</i> and <i>HIPPI</i> by siRNA technology	57
2.2.4	Microscopic technique	
2.2.4.1	Fluorescence microscopy	58
2.2.4.2	Confocal microscopy	58
2.2.5	Gene expression study	
2.2.5.1	RNA Isolation and first strand cDNA synthesis	58
2.2.5.2	Semiquantitative RT-PCR and Real Time PCR	59
2.2.5.3	Microarray study	59
2.2.5.4	Analysis of Microarray data	59
2.2.6	Sub-cellular fractionation and Western blot analysis	62
2.2.7	Immunoprecipitation assay	
2.2.7.1	Co-immunoprecipitation (CoIP)	63
2.2.7.2	Chromatin immunoprecipitation assay (ChIP) and Re-Chromatin immunoprecipitation (ReChIP) assay	63
2.2.8	Luciferase assay	65
2.2.9	Bioinformatics study	
2.2.9.1	Development of search tools	65
2.2.9.2	Transcription factor binding site analysis	66
2.2.9.3	Dataset preparation for Huntington's disease microarray analysis	66
2.2.10	Statistical analysis	67

Chapter 3: Results

3.1	Role of HIP1 in HIPPI mediated transcription regulation of <i>Caspase1</i> gene	69
3.1.1	HIP1 is necessary for HIPPI mediated transcriptional activation of <i>Caspase1</i> gene	69
3.1.2	Nuclear translocation of HIPPI is mediated by HIP1	70
3.1.3	Interaction of HIPPI-HIP1 heterodimer with the <i>Caspase1</i> gene promoter	73
3.1.4	Conclusion	74
3.2	Global role of HIPPI and HIP1 as transcription regulators: probable implications in HD pathogenesis	74
3.2.1	Genome wide search for the presence of HIPPI binding sites in the upstream of genes	74
3.2.2	Enrichment of other transcription factor binding sites in the vicinity of HIPPI binding site	75
3.2.3	Alteration of gene expression by HIPPI: microarray study	76
3.2.4	Validation of subset of genes altered in microarray study	80
3.2.5	Interaction of HIPPI with p53 in the nuclear compartment of cell and its effect on <i>Caspase1</i> expression	81
3.2.6	HIPPI binding sites at the deregulated genes in the caudate of HD patients: possible role of HIPPI in HD	83
3.2.7	Single gene analysis for HIPPI mediated transcription regulation in SHSY5Y cells	85
3.2.8	Conclusion	86
3.3	Regulation of RE1 Silencing Transcription Factor (REST) expression by HIPPI	86
3.3.1	Expression of RE1 silencing transcription factor (REST) is increased in presence of HIPPI in HeLa cells	87
3.3.2	HIPPI interacts with the upstream promoter of <i>REST</i> and increases its expression in Neuro2A cells	88
3.3.3	Effect of HIP1 and HIPPI mutations on <i>REST</i> and <i>BDNF</i> expression in Neuro2A cells	91
3.3.4	Knock down of <i>HIPPI</i> in Neuro2A cells results in <i>REST</i> down-regulation and <i>BDNF</i> up-regulation	93
3.3.5	Altered sub-cellular distribution of HIPPI and HIP1 in <i>STHdh</i> ^{Q111/111} cells, a cell model of HD	94
3.3.6	Expression of <i>REST</i> in <i>STHdh</i> ^{Q111/Q111} cells is higher compared to <i>STHdh</i> ^{Q7/Q7} cells	96
3.3.7	Knock down of HIPPI in <i>STHdh</i> ^{Q111/Q111} cells results in lowering of <i>REST</i> expression	98
3.3.8	Increased expression of <i>REST</i> in <i>STHdh</i> ^{Q111/Q111} cells is due to increased occupancy of HIPPI in <i>REST</i> promoter	98
3.3.9	Conclusion	99

Chapter 4: Discussion	102
4.1 The problem of nuclear localization of HIPPI: involvement of HIP1	103
4.2 HIPPI as global transcription regulator	104
4.3 Implication of HIPPI mediated transcription regulation in HD pathogenesis	109
4.4 Regulation of <i>REST</i> transcription by HIPPI in HD cell model	111
Summary and Conclusion	117
Appendix	121
Bibliography	139
List of Publication	163